

Management of Massive Transfusion Guidelines

Author and Contact details:	Practice Educator (Transfusion Lead). Tel: [REDACTED] Email: [REDACTED]	
Responsible Director:	Medical Director	
Approved by and date:	Clinical Effectiveness & Services Group	August 2018
Document Type:	TRANSFUSION GUIDELINE (Aintree Document)	Version 3.0
Scope:	All trust employees.	
Document Approval, History/Changes	For further information contact the Governance Department on Tel: [REDACTED]	

Think of the environment...Do you have to print this out this document? You can always view the most up to date version electronically on the Trust intranet.

1. Introduction

This guideline is to be followed for the management of all patients undergoing a massive blood transfusion.

2. Scope

All clinical and laboratory staff managing a patient with a massive blood transfusion.

This document is intended to be used in conjunction with “The Northwest Regional Toolkit for the Management of Massive Haemorrhage.”

3. Clinical activation of the guidelines

The Massive Haemorrhage guidelines will be activated for a patient requiring a massive blood transfusion. This can be defined as patients with:

- 1 Blood Volume loss per 24 hours
- Or > 50% blood volume loss in 3 hours
- Or ongoing blood losses > 150mls per hour

(Definition from The British Committee for Standards in Haematology)

The transfusion laboratory will not issue Major Haemorrhage Packs (MHP) for the *anticipation* of a massive haemorrhage for a patient in whom there is currently no active major bleeding as defined above. The only exception to this is in cases of **ruptured** abdominal aortic aneurysms who are to undergo surgery imminently, in which case one MHP may be ordered by a consultant only.

Where there is an anticipation of MHPs being required, and it is possible the Massive Haemorrhage Guidelines will be activated, it is advisable to alert the laboratory to this possibility at the earliest opportunity. This enables resources to be put in place in the laboratory to facilitate timely and expeditious issue of MHPs should the need then subsequently arise.


3.1. Documents for Use at The Walton Centre:

- Transfusion Management of Massive Haemorrhage in Adults
- Laboratory Algorithm
- “Seven Steps For Successful Coordination Of Massive Haemorrhage: Walton Centre / Aintree”
- Transfusion Management of Massive Haemorrhage in children (See Appendix 1)

4. References

“The Northwest Regional Toolkit for the Management of Massive Haemorrhage.”

Transfusion Management of Massive Haemorrhage in Adults



Thromboprophylaxis should be considered when patient stable

ABG – Arterial Blood Gas
FFP- Fresh Frozen plasma
PT- Prothrombin Time

APTT – Activated partial thromboplastin time
MHP – Massive Haemorrhage Pack
XM - Crossmatch

ATD- Adult Therapeutic Dose
NPT – Near Patient Testing

Laboratory Management of Massive Haemorrhage

Massive Haemorrhage Pathway Activated

Transfusion receives Call

'Massive Haemorrhage, Location, Specialty'
On standby

Receive call from designated communication lead in clinical area:

'This relates to massive haemorrhage situation'

The caller will state:

- Name and contact telephone number, name of consultant responsible
 - Patient's ID (surname, forename, hospital number, DOB or minimum acceptable patient identifiers if unknown)
 - Requirements:
 - State if O Neg is required as part of 1st pack (normally up to 2 packs)
 - Order massive haemorrhage pack 1
 - Clarify urgency of requirements to decide on need for further emergency group O, or time to wait for group specific or crossmatched red cells (issue as part of pack 1)
 - U+E, FBC, PT, APTT, Fibrinogen, ABG*, Adjusted Calcium, lactate*
- *may be near patient test

Receive samples and request forms

Haematology

Perform FBC, PT, APTT, Fibrinogen

Ring results to communication lead when available

Receive further calls from communication lead in clinical area:

Repeat investigations
Order for MHP 2
Liaise with on call haematologist (consultant / SpR)
Order for further components dependent on ongoing results
Stand down

Transfusion

Perform Group, antibody screen and crossmatch

Prepare MHP 1

Red cells* 4 units
(*emergency group O blood, group specific blood, XM'd blood depending on urgency)
FFP (group specific) 4 units
(lab staff to ensure stock platelets available or order 2ATD on blue light from blood centre)

Ring clinical area (communication lead) when blood / components ready

Prepare MHP 2

Red cells 4 units
FFP 4 units
Platelets 1 ATD
Cryoprecipitate 2Packs (if requested)

Restock Emergency Group O blood if used
Complete traceability audit trail

v1 2012

Seven Steps for Successful Coordination in Massive Haemorrhage: Aintree

1. Recognise trigger and activate pathway for management of massive haemorrhage; assemble the emergency response team

Phone [REDACTED] and request Outreach (if applicable).
Call consultant responsible for care of patient (if out-of-hours on call cons)

2. Allocate team roles

- i. Team leader
- ii. Communication lead– dedicated person for communication with other teams, especially the transfusion laboratory and support services
- iii. Sample taker / investigation organiser / documenter
- iv. Transporter - HCA, or other transfusion-trained member of team from clinical area)

3. Complete request forms / take blood samples, label samples correctly / recheck labelling

U+E, FBC, Crossmatch, PT, APTT, Fibrinogen, ABG, Calcium, lactate

Form should be an addressograph (preferable) otherwise handwritten with 4 identifiers (full name, DOB, hosp/NHS No)

Sample: Pink EDTA tube **must be handwritten**, (full name, DOB, hosp/NHS No) if emergency unknown patient then 2 identifiers are accepted (e.g. 'unknown 006785').

4. Request blood / blood components

Team leader should decide on use of:

- i. Emergency O Neg (immediate)

O neg blood available from transfusion laboratory (5 mins).
2 units O neg blood available in Recovery Fridge

- ii. Group specific **20 mins from receipt of sample**
- iii. Full Crossmatch **45 mins from receipt of sample**

Communication lead to contact laboratory:

Hotline: ☎ [REDACTED] or bleep [REDACTED] out-of-hours if no response on hotline

and inform the BMS of the following:

- i. Your name, location and ext number
- ii. 'this relates to the massive haemorrhage situation'
- iii. The patient's details: ideally surname, forename, hospital number, DOB (if unknown casualty: the hospital number and the unknown person number)
- iv. Whether O Neg will be required and how many units (usually 2)
- v. Order massive haemorrhage pack(s)
- vi. Contact lab if blood has been transferred with patient from another Trust (transferred blood must be sent in a box to the laboratory at Aintree prior to use) or patient is being transferred to another Trust

5. The clinical / laboratory interface

- i. Communication lead to arrange for transport of samples / request form to the laboratory
- ii. BMS to ring communication lead with results of urgent investigations
- iii. BMS to ring communication lead when blood / blood components are ready
- iv. Communication lead to arrange to collect blood and blood components from the laboratory

Designate HCA or other suitable staff member to transport samples to the laboratory.
The HCA must alert laboratory reception staff of the urgency of the request when delivered.

6. Communicate stand down of pathway and let lab know which products have been used

7. Ensure documentation is complete

- i. Clinical area: monitoring of vital signs, timings of blood samples and communications (lab and cons haematologist), transfusion documentation in patient casenote record, return traceability information to laboratory.
- ii. Laboratory: keep record of communications / telephone requests in patient laboratory record

Appendix 1 - Transfusion Management of Massive Haemorrhage in children

Transfusion Management of Massive Haemorrhage in Children

Ensure a consultant is aware of the massive haemorrhage and a senior member of staff is available to take charge of resuscitation if not already present.

Ongoing severe bleeding (overt / covert) and received 20ml/kg of red cells or 40ml/kg of any resuscitation in preceding hour.
Signs of hypovolaemic shock and / or coagulopathy
Administer Tranexamic acid (especially in trauma – ideally within 1 hour)
15mg/kg bolus over 10 mins (max 1000mg) intravenously: then infuse 2mg/kg/hr (max 125mg/hr) intravenously until bleeding is controlled.

Aintree University Hospitals NHS Foundation Trust
North West Regional Transfusion Committee
Incorporating North Wales

Activation Tel Number
Hotline [redacted]
Bleep [redacted] outside core hours if no reply on hotline
Emergency O red cells*
- location of supply:
Collect from Transfusion Lab
*Estimated time to receive at this clinical area: 5 min

- Group specific red cells: 20 min from receipt of sample
- XM red cells: 45 min from receipt of sample

Transfusion lab [redacted] or [redacted]
Hotline [redacted] or [redacted]
Consultant Haematologist via switchboard

Activate Massive Haemorrhage Pathway

RESUSCITATE
Airway
Breathing
Circulation

Call for help
'Massive Haemorrhage, Location, Specialty'
Alert: Outreach (if appropriate), on call consultant and patient's consultant, blood transfusion laboratory.
Consultant involvement is essential

Continuous cardiac monitoring

Take bloods and send to lab:
XM, FBC, PT, APTT, fibrinogen, U+E, Adjusted Ca
NPT: (A)BG, TEG / Rotem if available
and
Order Massive Haemorrhage Pack 1

Prevent Hypothermia
Use fluid warming device
Used forced air warming blanket

STOP THE BLEEDING

Give MHP 1
Red cells and FFP: give 10ml/kg in aliquots in a 1:1 ratio, reassess blood loss and respond to treatment; repeat as necessary

Consider 0.2 ml/kg 10% calcium chloride (max 10ml) over 30 mins

Further cryoprecipitate (10ml/kg) if fibrinogen < 1.5g/l or as guided by TEG / Rotem

Haemorrhage Control
Direct pressure / tourniquet if appropriate
Stabilise fractures
Surgical intervention (consider damage limitation surgery)
Interventional radiology
Endoscopic techniques

Reassess
Suspected continuing haemorrhage requiring further transfusion
Take bloods and send to lab:
FBC, PT, APTT, fibrinogen, U+E, Adjusted Ca (A) BG
Order MHP 2 (see table 2)
When half of MHP1 has been used consider ordering MHP2

Aims for therapy
Aim for:
Hb 8-10g/dl
Platelets >75 x 10⁹/l
PT ratio < 1.5
APTT ratio < 1.5
Fibrinogen >1.5g/l
Adjusted Ca >2.0 mmol/l
Temp > 36°C
pH > 7.35 (on ABG)
pH > 7.25 (on ABG)
Monitor for hyperkalaemia

Haemostatic Drugs
Vit K and Prothrombin complex concentrate for warfarinised patients
Other haemostatic agents: discuss with Consultant Haematologist

Give MHP 2
Red cells and FFP: give 10ml/kg in aliquots in a 1:1 ratio, reassess blood loss and respond to treatment; repeat as necessary.
Platelets: give up to 10ml/kg
Cryoprecipitate: give up to 10ml/kg

Thromboprophylaxis should be considered when patient stable

STAND DOWN
Inform lab
Return unused components
Complete traceability tags & documentation
Including audit proforma

Once MHP 2 administered, repeat bloods:
FBC, PT, APTT, fibrinogen, U+E, Adjusted Ca
NPT: (A) BG
To inform further blood component requesting

ABG – Arterial Blood Gas
APTT – Activated partial thromboplastin time
FFP- Fresh Frozen plasma
MHP – Massive Haemorrhage Pack
NPT – Near Patient Testing
PT- Prothrombin Time
XM - Crossmatch